

# GET TO KNOW YOUR SureClick® AUTOINJECTOR

When you're prescribed a medication with an Autoinjector, you might not know what to expect with self-administration. The ready-to-use SureClick® Autoinjector was designed for simple at-home use. This resource may help with questions about using the SureClick® Autoinjector.


## Important:

This resource is intended to provide you with an overview and basic information about the SureClick® Autoinjector and does not replace the Instructions for Use. Please review the Instructions for Use included with your prescribed medication with your doctor prior to performing your injection. If you have further questions please contact your doctor.



The images of the autoinjector are for illustrative purposes. The color of the autoinjector components varies based on the dose.


## KEY INJECTION STEPS



### PUSH

the Autoinjector firmly down onto the skin until the safety guard stops moving.

**This unlocks the start button.**



### PRESS

the start button; you will hear a **CLICK**. **KEEP HOLDING** the Autoinjector down.



### WATCH & CONFIRM

the window is fully **YELLOW**. It is okay to lift your thumb. It takes about 15 seconds to complete. You may hear a second click.

**Then you're done!**

**⚠ REMEMBER:** You **MUST** unlock the autoinjector by firmly pushing into your skin **WHILE** pressing the start button to perform the injection.

## APPROVED USE

Aimovig® (erenumab-aooe) is a prescription medicine used for the preventive treatment of migraine in adults.

## IMPORTANT SAFETY INFORMATION

### Who should not use Aimovig®?

Do not use Aimovig® if you are allergic to erenumab-aooe or any ingredients in Aimovig®.

**Before starting Aimovig®,** tell your healthcare provider (HCP) about all your medical conditions, including if you are pregnant or plan to become pregnant, breastfeeding or plan to breastfeed.

Tell your HCP about all the medicines you take, including any prescription and over-the-counter medicines, vitamins, or herbal supplements.

**Please see additional Important Safety Information throughout.**



**SureClick®  
Autoinjector**

# INJECTION RESOURCES FOR YOU

Are you getting started at home? Feel confident in knowing you have supplemental injection support online, on the phone, or at home.



## Instructions for Use

You can find the Instructions for Use at [aimovig.com/demo](https://aimovig.com/demo)



## Injection Overview Video

Hold your smartphone camera over the QR code to access the overview video, or find it at [aimovig.com/demo](https://aimovig.com/demo)



## Website

Online support is available to you. Visit [aimovig.com/demo](https://aimovig.com/demo)



## Live Support

We're here for one-on-one live support.  
Call us at 1-833-AIMOVIG (1-833-246-6844) 8 AM–9 PM ET, Mon–Fri



## Help From Your Doctor

If you have further questions, please contact your doctor.  
Call or email questions to:

NAME

PHONE

EMAIL



## Important Safety Information (cont'd)

### What are possible side effects of Aimovig®?

#### Aimovig® may cause serious side effects, including:

- **Allergic reactions.** Allergic reactions, including rash or swelling can happen after receiving Aimovig®. This can happen within hours to days after using Aimovig®. Call your HCP or get emergency medical help right away if you have any of the following symptoms of an allergic reaction: swelling of the face, mouth, tongue or throat, or trouble breathing.
- **Constipation with serious complications.** Severe constipation can happen after receiving Aimovig®. In some cases people have been hospitalized or needed surgery. Contact your HCP if you have severe constipation or constipation associated with symptoms such as severe or constant belly pain, vomiting, swelling of belly or bloating.

- **High blood pressure.** High blood pressure or worsening of high blood pressure can happen after receiving Aimovig®. Contact your healthcare provider if you have an increase in blood pressure.

The most common side effects of Aimovig® are pain, redness, or swelling at the injection site and constipation.

These are not all of the possible side effects of Aimovig®. Call your HCP for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](https://www.fda.gov/medwatch), or call 1-800-FDA-1088.

**Please see accompanying Aimovig® full Prescribing Information, including Patient Product Information.**



© 2023–2024 Amgen Inc.  
All rights reserved. Not for Reproduction. USA-334-85959

